

The kernel of the adjacency matrix of a rectangular mesh

Carlos Tomei and Tania Vieira *

Abstract

Given an $m \times n$ rectangular mesh, its adjacency matrix A , having only integer entries, may be interpreted as a map between vector spaces over an arbitrary field K . We describe the kernel of A : it is a direct sum of two natural subspaces whose dimensions are equal to $\lceil c/2 \rceil$ and $\lfloor c/2 \rfloor$, where $c = \gcd(m+1, n+1) - 1$. We show that there are bases to both vector spaces, with entries equal to 0, 1 and -1 . When $K = \mathbb{Z}/(2)$, the kernel elements of these subspaces are described by rectangular tilings of a special kind. As a corollary, we count the number of tilings of a rectangle of integer sides with a specified set of tiles.

1 Introduction

A basic combinatorial problem associated to a quadriculated surface is the counting of its tilings by dominos. Tilings of rectangles of integral sides, for example, were counted by Kasteleyn [K] in his solution of the Ising model. More generally, Kasteleyn showed how to slightly alter the adjacency matrix of a quadriculated disk so that the determinant of the resulting *Kasteleyn matrix* equals the desired count. Here, the adjacency matrix specifies which squares of the quadriculated surface share an edge. Tilings of other surfaces have been counted: the *Aztec diamond* is another example, presented in [EKLP]. Moreover, tilings have been *q-counted*, according to a height variable q ([EKLP] and [STCR]).

In [DT], the authors proved, in a different language, that the (-1) -counting of tilings of a quadriculated disk is always equal to 0, 1 or -1 . Their interest in the issue stemmed from the fact that the adjacency matrix A of a quadriculated surface is closely related to a standard discretization of the Laplacian with Dirichlet boundary conditions [C]. The result indicates a *spectral rigidity* for the discrete Laplacian, which so far has no interpretation in the continuous case. Further study of this rigidity led to consider additional properties of (-1) -counting, both on disks [STCR] and on annuli [ST1].

Since the adjacency matrix A has only integer entries, it may be interpreted as a linear transformation A_K between vector spaces over an arbitrary field K . By coloring squares of the quadriculated disk black and white in a chessboard fashion, it is easy to see that A_K takes vectors supported on the set of black squares (forming a subspace B_K) to vectors supported on white

*Departamento de Matemática, PUC-Rio; tomei@mat.puc-rio.br, tvieira@mat.puc-rio.br.
MSC Classification: Primary 05B45; Secondary 05C50.

squares (in W_K), and vice-versa. Thus, A_K splits naturally into two linear maps, $BW_K : B_K \rightarrow W_K$ and $WB_K : W_K \rightarrow B_K$ and the kernel of A_K is the direct sum of the kernels of BW_K and WB_K .

In this paper, we describe the kernel of A_K in the special case of an $m \times n$ rectangle quadrilaterated by (mn) squares of side of length 1. Suppose that the left lower square is painted black. As usual, the standard roundoff operations are denoted by $\lceil x \rceil$ and $\lfloor x \rfloor$.

Theorem 1. *For an $m \times n$ rectangle quadrilaterated by unit squares, $\dim \ker BW_K = \lceil c/2 \rceil$ and $\dim \ker WB_K = \lfloor c/2 \rfloor$, where $c = \gcd(m+1, n+1) - 1$. There is a basis to each kernel for which all vector coordinates are equal to 0, 1 or -1 .*

More recently [ST2], it was shown that, for an arbitrary quadrilaterated disk, both matrices BW_K and WB_K admit LDU -type decompositions, where the matrices L , D and U have all entries equal to 0, 1 or -1 . From this result, the fact that $\dim \ker BW_K$ and $\dim \ker WB_K$ are independent of K and the existence of bases for the kernels with entries equal to 0, 1 and -1 follows. The decomposition also implies the rigidity result in [DT].

When $K = \mathbb{Z}/(2)$, the discretization of the Laplacian essentially coincides with the adjacency matrix $A_{\mathbb{Z}/(2)}$ and we call the elements on $\ker BW$ and $\ker WB$ the *polarized $\mathbb{Z}/(2)$ -harmonic* functions, supported respectively on black and white points. Such objects have already been considered in [DT] as a technical tool. Draw the $m \times n$ rectangle so that the integer points of the auxiliary rectangle $R = [0, m-1] \times [0, n-1] \subset \mathbb{R}^2$ correspond to the centers of the squares. Consider tilings of R by tiles given in figure 1. Tiles can be rotated by integer multiples of $\pi/2$ and their dashed sides ought to belong to ∂R , the boundary of R . Length of sides is indicated in the picture, angles must be $\pi/4, \pi/2$ or $3\pi/4$. Figure 2.1 shows a tiling of the 7×4 rectangle.




Figure 1: The tiles; a and b are integers.




Figure 2: A tiling and a polarized $\mathbb{Z}/(2)$ -harmonic function of the 7×4 rectangle.

Integer points of R inherit the coloration of the squares (in particular, the origin is black). As we shall see in Proposition 2, tilings of R are such that the vertices of all tiles have integer coordinates and, with the possible exception of the corners of R , all vertices have the same color. Thus, the set of all tilings $\mathcal{T}$ naturally splits into two sets, $\mathcal{T}_b$ and $\mathcal{T}_w$. Polarized $\mathbb{Z}/(2)$ -harmonic functions are visually very simple: this is the content of the theorem below.

Theorem 2. *There are bijections*

$$\Phi_b : \ker BW_{\mathbb{Z}/(2)} \setminus \{0\} \rightarrow \mathcal{T}_b \text{ and } \Phi_w : \ker WB_{\mathbb{Z}/(2)} \setminus \{0\} \rightarrow \mathcal{T}_w$$

Combining both theorems, we obtain the number of elements in $\mathcal{T}_b$ and $\mathcal{T}_w$. Also, given a tiling (i.e., a polarized $\mathbb{Z}/(2)$ -harmonic function), there is a *lifting* to an element of $\ker BW_K \cup \ker WB_K$, having only 0, 1 and -1 as entries. This correspondence is essentially bijective: lifting of a vector can take at most two values v and $-v$. This is the content of Theorem 4. Lifting first appeared in [DT].

The first author acknowledges support from CNPq and FAPERJ, the second acknowledges support from CNPq. Both are thankful to Nicolau Saldanha for many conversations.

2 Kernel vectors

As before, let $R = [0, m-1] \times [0, n-1] \subset \mathbb{R}^2$. Adjacency in the $m \times n$ rectangular graph $G = R \cap (\mathbb{Z} \times \mathbb{Z})$ is the natural one: (x, y) and (x', y') are adjacent if and only if $x = x', y = y' \pm 1$ or $x = x' \pm 1, y = y'$. We call *points* the elements of G : *vertices* will be the corners of tiles. Color a point $(x, y) \in G$ *black* (resp. *white*) if $x + y$ is even (resp. odd): thus, the origin is black. Number black and white points in G from 1 to n_b and from 1 to n_w . Set BW to be the $n_w \times n_b$ *black-to-white adjacency matrix*: $BW_{wb} = 1$ if point b is adjacent to point w ; otherwise, $BW_{wb} = 0$. Similarly, define WB , the $n_b \times n_w$ *white-to-black adjacency matrix* of G . Since both matrices BW and WB only have integer entries, they may be interpreted as linear transformations between vector spaces over an arbitrary field K .

We compute the dimensions of the kernels of BW and WB and show they are independent of the base field K . The kernel elements admit two special symmetries, which we now describe. Given a vector v belonging to K^{mn} , denote by $\sigma_{i,j}$ the sum of the coordinates of v at the four neighbors of (i, j) in G , $\sigma_{i,j} = v_{i,j-1} + v_{i,j+1} + v_{i-1,j} + v_{i+1,j}$. Here $v_{i,j} = 0$ if $(i, j) \notin G$. Recall that the kernel of the adjacency matrix A of G is the direct sum of the kernels of BW and WB . Clearly, $v \in \ker A$ if and only if $\sigma(i, j) = 0$ for all points $(i, j) \in G$. Without loss, the sides of the rectangular graph satisfy $m \geq n$.

Proposition 1. *Let $v \in \ker A$. Then*

- (i) $v_{i,j} = v_{j,i}$ (resp. $-v_{j,i}$) if both points belong to G and $i + j$ is even (resp. odd).
- (ii) If v equals zero along the column $x = m_0$ then $v_{m_0-i,j} = -v_{m_0+i,j}$, if both points belong to G .

There are analogous symmetries across the other diagonals of the graph, and across rows of zeros of v .

Proof. Compare values of v along the lines $y = x + 1$ and $y = x - 1$. Since $\sigma_{0,0} = 0$, we must have $v_{1,0} = -v_{0,1}$. Sequentially, $\sigma_{1,1} = 0$, $\sigma_{2,2} = 0, \dots$ imply in turn $v_{2,1} = -v_{1,2}$, $v_{3,2} = -v_{2,3}, \dots$. Compare now values of v along the lines $y = x + 3$ and $y = x - 3$: making use of $\sigma_{2,0} = 0 = \sigma_{0,2}$, we have $v_{0,3} = -v_{3,0}$, $v_{1,4} = -v_{4,1}, \dots$. Proceed until the lines $y = x \pm (2k + 1)$ cover the square $[0, n - 1] \times [0, n - 1]$: this proves the statement for coordinates (i, j) for which $i + j$ is odd. The statement for coordinates with $i + j$ even follows from a similar sequential computation along lines $y = x \pm 2k$: to start off the induction argument, notice that $v_{i,j} = v_{j,i}$ along the diagonal $y = x$.

The proof of (ii) is a simple induction on i . Case $i = 1$ is as hard as the general case, and follows from expanding $0 = \sigma_{m_0,j} = v_{m_0-1,j} + v_{m_0+1,j}$. $\square$

Consider the $m \times n$ rectangular graph G and let $c = \gcd(m+1, n+1) - 1$. The *grid* $\mathcal{G}$ of G is the set of points in G belonging to the union of the horizontal lines $y = k(c+1) - 1$ with the vertical lines $x = k(c+1) - 1$, for integer k . The graph G , after removal of its grid $\mathcal{G}$, splits into $c \times c$ squares, called *fundamental squares*. Figure 3 shows the grid and the fundamental squares for the 14×9 rectangular graph. The figure also contains the description of an element v in $\ker WB$, interpreted as a vector with coordinates over the field $\mathbb{Z}/(2)$ of two elements — squares indicate the coordinates with nonzero entries. The edges connecting points in G are drawn to provide a visual indication of the symmetries of this kernel element. As we shall prove later, $\mathbb{Z}/(2)$ —polarized harmonic functions correspond to tilings of R : figure 3 is indeed the tiling associated to v .




Figure 3: Grid, fundamental squares and symmetries.

As before, fix an arbitrary base field K .

Theorem 3. *Let G be the $m \times n$ rectangular graph with grid $\mathcal{G}$ and adjacency matrices $BW_{m,n}$ and $WB_{m,n}$. Let $v \in \ker BW_{m,n}$ (resp. $\ker WB_{m,n}$). Then*

- (i) *The vector v is identically zero on $\mathcal{G}$.*
- (ii) *The restriction of v to each fundamental square lies in the kernel of the matrix $BW_{c,c}$ (resp. $WB_{c,c}$) of this square.*

(iii) Let s_1 and s_2 be two adjacent fundamental squares, bounded by a common row or column ℓ of $\mathcal{G}$. Then the restriction of v to s_2 is the odd extension of the restriction of v to s_1 across ℓ .

(iv) Let $G_{c,c}$ be the $c \times c$ square graph. Then

$$2 \dim \ker BW_{c,c} = c + (c \bmod 2) \quad \text{and} \quad 2 \dim \ker WB_{c,c} = c - (c \bmod 2).$$

(v) $\dim \ker BW_{m,n} = \dim \ker BW_{c,c}$ and $\dim \ker WB_{m,n} = \dim \ker WB_{c,c}$.

(vi) There are bases for $\ker BW_{m,n}$ and $\ker WB_{m,n}$ consisting of vectors having coordinates taking only the values 0, 1 and -1 .

In particular, the dimensions of $\ker BW_{m,n}$ and $\ker WB_{m,n}$ are independent of the base field K .

Proof. We shall prove (i), (ii), and (iii) by induction on the set of pairs (m, n) for which $\gcd(m+1, n+1) = c+1$ is fixed, ordered lexicographically. For the starting point $m = n = c$, the result is obvious. Suppose that $v^{m,n} \in \ker BW_{m,n}$, in a self-explanatory notation. From the symmetry (i) of the previous proposition, the restriction of $v^{m,n}$ to column $x = n$ is zero (recall that $m \geq n$). Indeed, for $v = v^{m,n}$, since

$$\begin{aligned} 0 &= \sigma_{n-1,j} = v_{n-2,j} + v_{n-1,j-1} + v_{n-1,j+1}, \\ 0 &= \sigma_{j,n-1} = v_{j,n-2} + v_{j-1,n-1} + v_{j+1,n-1} + v_{j,n}, \end{aligned}$$

we must have $v_{j,n} = 0$. Removal of column $x = n$ from $G = G_{m,n}$ gives rise to two connected subgraphs: a square $G_{n,n}$ containing the first n columns of G and a possibly rectangular subgraph $G_{m-n-1,n}$ containing only the last $m - n - 1$ columns. Also, by making use of the column of zeros and the fact that $v^{m,n} \in \ker BW_{m,n}$, we have that the restriction $v^{m-n-1,n}$ of $v^{m,n}$ to $G_{m-n-1,n}$ belongs to $\ker BW_{m-n-1,n}$. Now, $\gcd(m-n-1+1, n+1) = c+1$: by induction, then, $v^{m-n-1,n}$ has zeros on the grid of $G_{m-n-1,n}$. Thus, the nonzero entries of $v^{m-n-1,n}$ belong to $c \times c$ squares framed by points in which $v^{m-n-1,n}$ is zero. Now using symmetry (ii), mirroring these squares across columns $n, n-c-1, \dots$, we see that the square $G_{n,n}$ also splits in $c \times c$ squares with the same property. Conversely, taking odd extensions of a kernel element on the fundamental square across the grid obtains a kernel element on the full rectangle: this proves (v). A vector in $\ker BW_{c,c}$ (or in $\ker WB_{c,c}$) is clearly determined by its values along the first column $x = 0$ of $G_{c,c}$. The fact that any choice of values along this columns indeed gives rise to a kernel vector follows from the diagonal symmetry (i) of Proposition 1 — this proves (iv). A special choice of vectors along the first column of $G_{c,c}$ obtains basis of kernel vectors with coordinates 0, 1 and -1 . For $\ker BW_{c,c}$, they are $e_1, e_3 - e_1, e_5 - e_3, \dots, e_{c+(c \bmod 2)-1} - e_{c+(c \bmod 2)-3}$ and for $\ker WB_{c,c}$, $e_2, e_4 - e_2, \dots, e_{c+(c \bmod 2)-2} - e_{c+(c \bmod 2)-4}$. The details are left to the reader: figure 4 describes the simple bases for $G_{7,7}$. Squares and circles correspond to 1's and -1 's respectively.

□

4 Defining the bijections Φ_b and Φ_w

For now on, the base field K is $\mathbb{Z}/(2)$. The rest of the paper is dedicated to the proof of the theorem 2.

Let G be the rectangular graph associated to R and take $u \in \ker BW \setminus \{0\}$. A black point $b \in G$ is an *active point of u* if $u(b) = 1$. Denote by A_u the set of active points of u . In A_u , consider the following *adjacency relation*. Two points are adjacent if both conditions are satisfied: (i) both points have a common white neighbor w in G ; (ii) both points form a right angle with w or they are the only active neighbors of w .




Figure 5: Adjacencies in A_u .

A *splitting* of R is a decomposition in *chunks*, i.e., polygons with integral vertices, which are not necessarily tiles. Given a vector $u \in \ker BW \setminus \{0\}$, let $\Phi_b(u) = \text{int}(R) \setminus \{\overline{b_1 b_2} \mid b_1 \text{ and } b_2 \text{ are adjacent in } A_u\}$. Said differently, draw straight line segments joining adjacent points in A_u . In Section 4.1, we will see that this construction obtains a splitting $\Phi_b(u)$ of R . In Section 4.3, more will be proved: $\Phi_b(u)$ is actually a tiling, i.e., its chunks are tiles.

4.1 Geometry of A_u

Let $b \in A_u$ be a black point in the interior of R . The *star* centered at b is the set of points in A_u which are adjacent to b , together with the segments joining b to these points. The Lemma below follows by exhausting possibilities.

Lemma 1. *Let $b \in A_u$ be an interior point of G . Up to rotations and reflections, the stars centered at b are listed in figure 6.*

We point out that all stars in the list indeed occur in tilings.

Lemma 2. *Every point of A_u which is not a corner of R has (at least) two neighbors in A_u . Also, segments $\overline{b_1 b_2}$ and $\overline{b_3 b_4}$ in A_u joining four distinct points do not intercept.*

Proof. Let b be a point in A_u which is not a corner of R . Then b has two neighbors in G , w_1 and w_2 , so that the three points are collinear. Since b is an active neighbor of w_1 and w_2 and $u \in \ker BW$, w_1 and w_2 have other active black neighbors in G . Let b_1 (resp. b_2) be the active neighbor of w_1 (resp. w_2) closest to b . Notice that b_1 and b_2 are distinct, since b is the only black neighbor common to w_1 and w_2 . Thus, b is adjacent in A_u to at least two points b_1 and b_2 .

We now consider the second statement. From the definition of the adjacency in A_u , the possible segments joining active points are given in figure 7. Since the four points are distinct, segments may not meet at endpoints. Also, as all points have the same color, the only plausible intersection of segments are listed in figure 7. By direct inspection, each such intersection violates the definition of adjacency in A_u . $\square$




Figure 6: Stars centered at b .




Figure 7: Segments and their possible intersections.

Proposition 3. *The closure of the connected components of $\Phi_b(u)$ are convex polygons with vertices in A_u or in the corners of R . In particular, $\Phi_b(u)$ is indeed a splitting.*

Proof. From Lemma 2, the juxtaposition of segments in A_u , together with ∂R , consists of closed polygonal curves and the vertices of the connected components in the statement are either in A_u or in the corners of R . From the classification of stars, convexity follows at interior vertices: for boundary vertices, convexity is trivial. $\square$

4.2 The possible sides and angles of chunks of $\Phi_b(u)$

Sides of chunks of a splitting $\Phi_b(u)$ are *active* if they consist only of segments of A_u . *Inactive sides* are those which contain no segment of A_u . The remaining sides are called *mixed*. Inactive and mixed sides must lie in ∂R , since sides in $\Phi_b(u)$ are union of segments in A_u with segments in ∂R .

Lemma 3. *There are no mixed sides.*

Proof. Suppose e is a mixed side of a chunk in $\Phi_b(u)$. Without loss, suppose e belongs to the upper side of R . The $m \times n$ rectangular graph G ought to include points immediately below e , since $m, n > 1$. Let b be a point of e joining an active and an inactive segment, and let w denote the white point below b . Figure 8 shows the possible active neighbors of w : any possible choice




Figure 8: Mixed segments do not exist; dashed segments are inactive.

implies the existence of a segment in A_u meeting e at b : in particular, e may not be a side of a chunk. $\square$

Lemma 4. *Horizontal or vertical active sides of chunks in $\Phi_b(u)$ have length 2.*

Proof. Consider a horizontal active side h of a chunk of $\Phi_b(u)$. From the adjacency relation in A_u , h has even length. Suppose h has length greater to 2, and take e in h joining two active points at distance 4.




Figure 9: Horizontal sides.

As in the previous lemma, we may suppose that the integral points immediately under h belong to G . The midpoint of e has at least one white neighbor $w \in G$, as shown in figure 9. From the adjacency relation in A_u , the side neighbors of w are inactive. There is only one possibility: the other active neighbor of w must be b , as in figure 9. But then, the vertical segment in the same figure ought to be a segment in A_u , contradicting the fact that e lies in a side of a chunk. $\square$

The angles of chunks are formed by the segments listed in figure 7 combined with segments in ∂R . Our next step will be the classification of angles according to activity of their sides. Angles defined by an active and an inactive side are *mixed angles*. From the adjacency relation in A_u , angles of $\pi/4$ in $\Phi_b(u)$ are mixed and angles of $3\pi/4$ are not. We are left with considering right angles.

Lemma 5. *Right angles are not mixed.*




Figure 10: An inexistent right angle.

Proof. Consider first the case in which the sides of the angle are horizontal and vertical, as in figure 10. The inactive side must then belong to ∂R . The number

of active neighbors of w must be even, and hence $u(b) = 1$. But then, the adjacency relation in A_u implies the presence of active segments as in figure 10.

The other kind of right angle has diagonal sides: these sides may not belong to ∂R and thus have to be active. Indeed, by the adjacency relation in A_u , this kind of right angle may only occur in square chunks of sides with length $\sqrt{2}$. $\square$

Adding up, figure 11 lists all possible angles of chunks in $\Phi_b(u)$, taking into account the activity of each side.




Figure 11: Angles; active and inactive sides.

Lemma 6. *The number of $\pi/4$ angles in a chunk of a splitting $\Phi_b(u)$ is even.*

Proof. Lemma 3 excludes the possibility of mixed sides. Clearly, every chunk ought to have an even number of mixed angles, and the only mixed angles measure $\pi/4$. $\square$

Lemma 7. *Let C be a chunk in $\Phi_b(u)$ with a side s common to two right angles which is active and either horizontal or vertical. Then C is a square with sides of length 2.*




Figure 12: (1) implies (2).

Proof. Lemma 4 implies that the side s has length 2. In figure 12.1, b is not active, again in accordance with the adjacency relation in A_u . Thus, the only active neighbors of w are adjacent in A_u . $\square$

4.3 Chunks in $\Phi_b(u)$ are tiles

Since the smallest external angle of a chunk is $\pi/4$ (figure 11) chunks of $\Phi_b(u)$ have at most 8 sides — we now classify chunks.

Octagons: All inner angles have to measure $3\pi/4$. Horizontal and vertical sides have length 2 since they are active (Lemma 4) and thus octagons must be as in figure 1.

There are no heptagonal chunks: A heptagonal chunk should have 6 angles measuring $3\pi/4$ and one, $\pi/2$ (figure 13). Horizontal and vertical sides must be active and have length 2. From figure 13, adding coordinates on both axis,

$a' - b' - c' = 0$ and $-a' - b' + c' = 0$. Thus, $a - b - c = 0$ and $-a - b + c = 0$, and hence $b = 0$: one side is gone.




Figure 13: There are no heptagonal chunks: $a = \sqrt{2}a'$, $b = \sqrt{2}b'$ and $c = \sqrt{2}c'$.

Hexagons: From Lemma 6, chunks of $\Phi_b(u)$ have an even number of angles measuring $\pi/4$. Thus, hexagons must have four $3\pi/4$ angles and two $\pi/2$ angles. Angles ought to be ordered as in the figure 1: from Lemma 7, the right angles have to be separated.

There are no pentagonal chunks: Again, from Lemma 6, a pentagon should have two $3\pi/4$ angles and three right angles. But then two consecutive right angles are inevitable, contradicting Lemma 7. Notice that the hypothesis of the Lemma is fulfilled: right angles are not mixed and hence no inactive sides.

Quadrilaterals: There are two cases. First consider rectangular chunks. At least one of its sides must be active, otherwise $u = 0$. Since right angles are not mixed, all sides must be active. But then, from Lemma 4 and the adjacency relation in A_u , the chunk must be one of the squares in figure 1. Second, the chunk may have two $3\pi/4$ angles and two $\pi/4$ angles and must be the trapezoidal tile in figure 1.

Triangles: The triangle must be right isosceles (figure 1).

Trapezoidal and triangular chunks have dashed sides: $\pi/4$ angles are mixed. The upshot of the classification is the following — *the possible chunks in a splitting $\Phi_b(u)$ are the tiles in figure 1: $\Phi_b(u)$ is a tiling.* Thus, the function Φ_b indeed takes nonzero vectors in the kernel of BW to the subset $\mathcal{T}_b$ of tilings. There is an analogous function Φ_w from $\ker WB \setminus \{0\}$ to $\mathcal{T}_w$.

5 Φ_b is a bijection

We start showing that $\Phi_b : \ker BW \setminus \{0\} \rightarrow \mathcal{T}_b$ is injective. Take $u, v \in \ker BW$, inducing tilings $\Phi_b(u)$ and $\Phi_b(v)$. We have to show that if $\Phi_b(u) = \Phi_b(v)$, then $A_u = A_v$ (and then, trivially, $u = v$). If $\Phi_b(u) = \Phi_b(v)$, then A_u and A_v may only differ on ∂R , since the only sides which may be active or not lie there. It is the shape of the tile which determines which side is active or not: inactive sides are the large basis of trapezoidal tiles and short sides of right triangles. Thus, sides of a tile in $\Phi_b(u) = \Phi_b(v)$ belonging to ∂R must be simultaneously active or not in both u and v , and the proof of injectivity is complete.

We now consider the surjectivity of Φ_b . Given a tiling $T \in \mathcal{T}_b$ (and in particular, the description of type of activity for each side), we have to find a nonzero vector $u \in \ker BW$ for which $\Phi_b(u) = T$. The sequence of steps below is natural.

- Defining u from T .
 $u(b) = 1 \Leftrightarrow b$ belongs to a non-dashed side in T .
- Checking that $u \in \ker BW$.

We must show that, for each white point w , the sum of the values $u(b)$ for its neighbors $b \in G$ is even. By checking for each kind of tile, it is clear that the sum of $u(b)$ for neighbors b of w in a single tile is even. Thus, if w lies in the interior of a tile, we are done. Suppose now that w lies in a side of a tile of T . First, notice that vertices of tiles belonging to active sides are black. Hence, if w is such a vertex, it must belong to inactive sides, and thus, both sides containing w lie in ∂R — w must be a corner of R , belonging to a right triangle. In this case, the black neighbors of w are active, if the short sides of the triangle are of length 1, or inactive, for longer lengths. In both cases, the total sum of $u(b)$ over black neighbors b of w is even.

Finally, if w belongs to a side of a tile without being a vertex, this side must be either horizontal or vertical, since diagonal sides only cross black points. If this side lies in ∂R , all the black neighbors of w belong to the same tile, and we are done. If, instead, this side is in the interior of R , we may take it to be horizontal and it must be an active side. Thus, $u(b_1) = u(b_2) = 1$, where b_1 and b_2 are the side neighbors of w . Besides, the black neighbors of w belong to two tiles, since w is not a vertex. Suppose b_1 , b_2 and b_3 in one tile and b_1 , b_2 and b_4 in another. Then $u(b_1) + u(b_2) + u(b_3)$ and $u(b_1) + u(b_2) + u(b_4)$ are even. Since $u(b_1) = u(b_2) = 1$, we must have $u(b_3) = u(b_4) = 0$. Hence, $u(b_1) + u(b_2) + u(b_3) + u(b_4) = 0 \pmod{2}$.

- The vector u obtained from T indeed satisfies $\Phi_b(u) = T$.

The black points by which non-dashed sides of T and $\Phi_b(u)$ passes are the same. Indeed, these are the active points of u . We are left with checking that the segments between these points are the same in T and $\Phi_b(u)$. Clearly, ∂R consists of segments common to both tilings. Therefore dashed sides are irrelevant: we only need to prove that T and $\Phi_b(u)$ have the same active segments.

The *active neighborhood* of a point $w \in G$ under $u \in \ker BW$ is the set of its active black neighbors in G . An active neighborhood is *trivial* if there are no active points. A simple check obtains all nontrivial active neighborhoods: they are listed in figure 14, up to rotations. In cases (1), (2) and (3), w is an interior point; in (4) and (5), w belongs to a side of ∂R , and in (6), w is a corner of R . Trivial active neighborhoods are not relevant in the forthcoming argument. In order to show that T and T_Φ have the same active segments, all we have to do is to show this fact for each active neighborhood.

In the case of a neighborhood of type (1), the two active points are joined by a segment in $\Phi_b(u)$, by definition. Suppose that this segment does not belong to T : w then is in the interior of a tile t of T . The two active neighbors of w belong to the boundary of t , by the construction of u from T . Since tiles are convex, the segments of T form angles which are less than or equal to π . Also, sides of tiles in T do not pass by points at which u is zero (we do not have to take into account boundary points). Thus, the sides of t passing by the active points of the neighborhood of w must be horizontal, as shown in figure 15; these sides extend to active neighbors at distance 2. But, by figure 1 horizontal sides have length at most 2 — contradiction. Thus T equals $\Phi_b(u)$, when restricted to an active neighborhood of type (1).

For a neighborhood of type (2), T contains active segments joining each




Figure 14: Nontrivial active neighborhoods.




Figure 15: The boundary of tile t .

neighbor of w to some other neighbor. Consider the possibility of one of this segments being horizontal, as $\overline{b_1 b_3}$ in figure 16.1. Neighbor b_2 is either joined to b_3 or to b_4 , up to trivial symmetries, as in figure 16. Tiling T cannot contain (1), since the $\pi/4$ angle may only appear at ∂R (again, check the shape of tiles in figure 1). Possibility (2) may not happen either, since it implies a white vertex of a tile which is not a corner of R . Thus segment $\overline{b_1 b_3}$ is not in T . Vertical segments are equally excluded.

Suppose now that there are only diagonal active segments. The only tile with parallel diagonal sides $\sqrt{2}$ apart is the tilted square. Thus, the four diagonal segments must appear in both T and $\Phi_b(u)$.




Figure 16: Active segments.

The study of neighborhoods of types (3), (4), (5) and (6) is similar and will be left to the reader. This proves the surjectivity of Φ_b .

At this point, the proof of Theorem 2 is complete.

A consequence of the theorem is the fact that tilings of the same color can be added, since they correspond to kernel elements of the same matrix. Figure 17 shows all tilings of the 10×4 rectangle: the nonzero elements in the kernel of WB give rise to the first three, A , B and $A+B$. The remaining ones correspond

to nonzero elements in the kernel of BW .




Figure 17: All tilings of the 10×4 rectangle.

We now consider the lifting of polarized $\mathbb{Z}/(2)$ -harmonic functions. Again, a tiling T corresponds to a vector $u \in \ker BW_{\mathbb{Z}/(2)} \cup \ker WB_{\mathbb{Z}/(2)}$. Consider its graph A_u of active points: from the shape of the tiles, each loop in the graph has even length. In particular, active points may be labeled 1 and -1 alternatively: this defines the *lifting* of u to a vector with coordinates equal to 0, 1 and -1 which may be interpreted as lying in $\ker BW_K \cup \ker WB_K$. The connectivity of A_u is all we need to prove the result below.

Theorem 4. *Lifting is a bijection between polarized $\mathbb{Z}/(2)$ -harmonic functions and classes of vectors, $[v, -v]$, where $v \in \ker BW_K \cup \ker WB_K$ with coordinates equal to 0, 1 and -1 .*

References

- [C] COLLATZ, L., The numerical treatment of differential equations, Springer-Verlag, Berlin (1966).
- [DT] DEIFT, P. A. & TOMEI, C., On the determinant of the adjacency matrix for a planar sublattice, *J. Combin. Theory Ser. B* **35**, no. 3, 278–289 (1983).
- [EKLP] ELKIES, N., KUPERPBERG, G., LARSEN, M., PROPP, J., Alternating sign matrices and domino tilings, *J. Alg. Combin.* **1**, 112–132 and 219–239 (1992).
- [K] KASTELEYN, P.W., The statistics of dimers on a lattice, I. The number of dimer arrangements on a quadratic lattice, *Physica* **27**, 1209–1225 (1961).
- [STCR] SALDANHA, N., TOMEI, C., CASARIN, M. AND ROMUALDO, D., Spaces of domino tilings, *Disc. Comp. Geom.* **14**, 207–233 (1995).
- [ST1] SALDANHA, N., TOMEI, C., Tilings of quadriculated annuli,
URL: <http://www.arxiv.org/math.CO/0012265>.
- [ST2] SALDANHA, N., TOMEI, C., Arithmetic properties of the adjacency matrix of quadriculated disks,
URL: <http://www.arxiv.org/math.CO/0107162>.